

1x RBC Lysis Buffer



Order No.	Size	Sterility
60-00050-11	100 ml	無菌状態
60-00050-12	250 ml	無菌状態
60-00050-13	500 ml	無菌状態

概要

Red Blood Cell (RBC) Lysis Buffer has been designed, formulated, and tested to ensure optimal lysis of RBCs in single cell suspensions with minimal effects on leukocytes. Nucleated RBCs are not effectively lysed with ammonium chloride. This 1x RBC Lysis Buffer is ready to use.

Components

- NH₄ Cl (155 mMol/L)
- KHCO₃ (10 mMol/L)
- EDTA (0.1 mMol/L)

Protocol

1. Ensure that the RBC Lysis Buffer is cold (4-8°C) .
2. Dilute one volume of cell suspension with 10 volumes of 1x Red Blood Cell Lysis Solution.
3. *When working with blood samples:*
Vortex for 5 seconds and incubate for 10-15 minutes (no more than 15 minutes) at 4-8°C.

When working with tissue samples:

Vortex for 5 seconds and incubate for 2 minutes at 4-8°C.

Note: Observe turbidity to evaluate red blood cell lysis. Once the sample becomes clear red, lysis is complete.

4. Vortex for 5 seconds and centrifuge at 300×g for 10 minutes at room temperature.
5. Aspirate supernatant completely.
6. Resuspend the cell pellet in an appropriate buffer and proceed to further applications.

Additional Information

納期

1-2

興味領域

細胞生物学

不妊症	無菌状態
フォーレンジ	pH 7.4
うんぱんじょうたい	室温
保管状態	4-8 °C
きせいぶん	For research use only. Not for use in diagnostic procedures.,

Warning and Limitations

This product is for research and development only, not for diagnostic or therapeutic use.